



Thomas Obermeyer, M.D.



obermeyermd.com

O: 847-285-4318

F: 847-885-0130



**BARRINGTON**  
Orthopedic Specialists  
*Specializing in You*

## POST-OPERATIVE INSTRUCTIONS: ORTHOPEDIC SURGERY

### MEDICATION

- A certain amount of pain, swelling and bruising is expected for the days following surgery. The goal of post-surgical pain medication is to *control* and *manage* post-surgical pain, not to *eliminate* it.
- Most patients will be given a prescription for a narcotic pain medication, Norco (hydrocodone) that is initiated shortly after surgery. Take it only as needed after your discharge.
  - Common side effects of the pain medication are nausea, drowsiness and constipation – to decrease the side effects take the medication with food. An over the counter stool softener (Colace/Ducolax) may be taken to prevent constipation.
  - You may not drive or operate heavy equipment while taking narcotic pain medicine.
  - Call the office immediately if you are having an adverse reaction to the medication.
- Resume all home medications unless otherwise instructed by Dr. Obermeyer or your primary care physician.
- You may be recommended to take a blood thinner, which is individualized to you based on risk for clotting after surgery. The type of medication is determined by Dr. Obermeyer and your surgical team. Commonly this medication is aspirin 81mg twice daily started the day after surgery and continued for a few weeks.

### WOUND CARE

- It is recommended you leave in place the bandage you left the surgical facility with until the five (5) day mark after surgery. It is recommended the incision remain clean under that bandage for those days to prevent infection and promote early healing. You may shower with a sealed bag over the bandages during that time, ensuring the bandages remain dry.
  - Once the bandages are removed, the incision can be left open to air or replaced with a gauze or ace wrap. Do not apply lotion, cream, or ointments to the incision.
  - Showering is permitted at five days with gentle soaping, and you can let water run over the staples and the front of the knee at that time. Please avoid *submerging* the knee in a bath for a few weeks after surgery to prevent contamination of the incision and infection.
- Mild bleeding and swelling is normal. If blood soaks onto the bandage do not become alarmed, reinforce with additional dressing.
  - If there is excessive drainage from the surgical incision requiring multiple reinforcements and dressing changes, please contact the office to inform Dr. Obermeyer and the surgical team.



Thomas Obermeyer, M.D.



**BARRINGTON**  
Orthopedic Specialists  
*Specializing in You*

- If there is excessive redness or drainage from the incision, or fever greater than 101.5°F, contact Dr. Obermeyer and the surgical team. Lower grade temperature elevations for a couple days after surgery are normal and should not be a cause for alarm.

#### COLD THERAPY

- Ice can be applied through a clean plastic bag, or a bag of frozen peas accomplishes the same objective. Formal icing machines are not covered by insurance and are not routinely used, but Dr. Obermeyer supports their use if you have access to one. **Never apply cooling agents (ice, polar care etc) directly to exposed skin.**
- After the first day, use 20 minutes on, 20 minutes off, as long as it continues to help with pain. You may discontinue when you feel the cooling is no longer helpful.
- Consider using ice after physical therapy to help with swelling and pain.

#### ACTIVITY

- Keep the involved limb elevated for several days to help with swelling.
- Depending on your limitations after surgery, several exercises will be beneficial for you at home: quad sets, ankle pumps and straight leg raises and should be done 3-4 times per day.
- Being up and around after surgery will help diminish the risk of blood clots.
- Formal physical therapy (PT) will typically begin within two weeks of surgery.

#### RETURN TO WORK

- Returning to work depends on the type of work you do. Patients in seated jobs may return to work in as little as one week if demands on you are limited and you have transportation arranged. You will possibly be taking narcotic pain medications and not have the ability to drive. Patients with higher demands may require a longer time frame and Dr. Obermeyer or his team will discuss this with you. Please let our office know if you need assistance with paperwork for your time away.

#### EMERGENCIES

- Contact Dr. Obermeyer's office (847-285-4200) if any of the following are present:
  - Uncontrolled nausea and vomiting
  - Uncomfortable or painful swelling that affects the surgical leg much more than the other normal one
  - Unrelenting pain at the knee or **pain in the calf**
  - Fever >101.5°
  - Redness or continuous drainage around incisions
- **If you are having chest pain or difficulty breathing, call 911 or go to the closest emergency room immediately for evaluation.**

#### FOLLOW-UP CARE



Thomas Obermeyer, M.D.



**BARRINGTON**  
Orthopedic Specialists  
*Specializing in You*

- Please make your first post-op visit 2 weeks after surgery if not done so already.
- **If you have any questions or concerns about the above please call Dr. Obermeyer and the team at 847-285-4318.**